

## Removal of sutures

Dear Doctor/Nurse,

Your Patient attended AllView Healthcare clinic today for the following procedure:

- Curettage
- Punch Biopsy
- Shave Biopsy
- Excision
- Mohs Surgery

**Sutures are due for removal on:**

Date: \_\_\_\_\_

Site: \_\_\_\_\_

\_\_\_\_\_

*Please note that we use a continuous suture for excisions & Mohs, this must be removed in sections to avoid disruption to the wound.*

Scan the QR code to download this leaflet, or to access the online version.



FM-AV-036a - V2.0 | Public

## What's next after your surgical procedure is completed



The sample taken during your surgical procedure is now **sent to the laboratory** for analysis by a histopathologist. If you have undergone Mohs Surgery with us, your results will be discussed with you on the day.



We expect to have the **results back in 4-6 weeks** post your procedure.



When we receive your results, **you will be contacted by our medical team** (GP or nurse) to discuss the results and treatment plan.

Signed by: \_\_\_\_\_


Print name: \_\_\_\_\_

# AllView Healthcare

Patient Information

## Postoperative Wound Care

for Dermatology Procedures



If you have any concerns, please contact a member of AllView Healthcare Surgical Nursing Team

**Phone: 01 224 8100**

**Email: [woundcare@allview.ie](mailto:woundcare@allview.ie)**

We are available between the hours:

**9 am – 5 pm Monday – Friday**

## After surgery

The nurse discharging you after your procedure will explain which type of wound closure was used and if you need to have them removed.

You may need to **attend your GP/Practice Nurse for removal of sutures**, contact them soon to ensure you get an appointment.

Your local anaesthetic can last 2 to 4 hours. If you feel some soreness or discomfort, a mild pain relief such as Paracetamol should be sufficient.

Ibuprofen or Aspirin are not recommended.

Alcohol and vitamin E tablets should be avoided for 48 hours after your operation as they can increase your risk of bleeding.

Please do not carry out heavy lifting or vigorous exercise as advised.

The skin around your wound may be numb for several weeks/months following surgery.

You may have some swelling and bruising around the wound. This can take over a week to settle down.

You may see a small amount of blood on your dressing – this is normal. **Please seek medical attention if the bleeding does not stop after 20 minutes** of continuous firm pressure.



### Signs of infection or postoperative complications:

- Increasing redness
- Excessive swelling
- Increasing or severe pain
- Increased temperature
- Generally feeling unwell
- Discharges pus

*Please contact us should you experience any of the above.*

**Phone: 01 224 8100**

**Email: [woundcare@allview.ie](mailto:woundcare@allview.ie)**

*If outside normal working hours, please contact your GP or present to your local Accident & Emergency Department.*

## Wound care

Your wound may be tender and bleed slightly when you change your dressings over the first few days following your surgery.

- Keep dressings dry and in place for 24/48 hours after your procedure.
- Afterwards change dressing daily or as needed/directed. **It is advisable to keep dressing on until the sutures have been removed.**
- Gently remove the dressing (it may be easier to get the dressing wet first).
- Clean the wound with cooled boiled water (cotton buds/pads may be used).
- Pat dry with tissue paper.
- Apply \_\_\_\_\_ to the wound to keep it moist.
- Cover the wound with a simple dressing.
- Continue these steps daily until the stitches are removed.
- Keep wound moist and greasy at all times, do not let it dry out or crust.

Initially your scar will be red and raised, this will reduce in colour and size over several months. **It can take 6 months for the scar to heal completely.**